

## THE GENETIC AND NON-GENETIC AFFECT ON CARCASS CHARACTERISTICS OF NEW ZEALAND WHITE AND BALADI BLACK RABBITS

التأثير الوراثي وغير الوراثي على خصائص الذبيحة في أرانب النيوزيلندي الأبيض والبلدي الأسود

By

Amira S.El-Deighadi; M. G. Gharib; M.I seif EL-Nasr;  
Yonan G. E. Mogda M. Salem and Lamia F. Abdel Mawla  
Animal Production Research Institute, Agriculture Research  
Center, Dokki, Cairo, Egypt (APRI).

Doi: 10.21608/asajs.2021.179614

قبول النشر: ٢٢ / ٦ / ٢٠٢١

استلام البحث: ٢٩ / ٥ / ٢٠٢١

El-Deighadi , Amira S.; Gharib , M. G.; seif EL-Nasr, M.I;  
Yonan, G. E. Salem, Mogda M. & Mawla ,Lamia F. Abdel  
(2021). *Arab Journal of Agriculture Sciences*,  
AIESA.(4),11, pp 141 - 162.

**ABSTRACT:**

Data collected on 120 random male at 12 weeks of age, from New Zealand white and Baladi Black rabbit. Data were analyzed applying multi-trait animal model of carcass characteristics to obtain proportion of the phenotypic variance due to additive genetic effects ( $h^2$ ); common litter effects ( $c^2$ ); random error effects ( $e^2$ ); genetic ( $r_G$ ), common litter ( $r_C$ ), environmental ( $r_E$ ) and phenotypic ( $r_P$ ) correlations. Data were carried out by weighed least-squares means method in the procedure GLM of statistical software SAS(2003) to obtain least squares of effects for breed, parity and season and significant. Heritability estimates were mostly moderate 0.20, 0.12 and 0.37 for live weight at slaughter, weight after bleeding and weight with viscera and without head, were moderate and high and ranged from 0.29 to 0.54 for edible parts, were moderate and ranged from 0.30 to 0.35 for dressing yield, also were low and moderate and ranged from 0.03 to 0.33 for inedible parts. Common litter effects Estimates were 0.37, 0.55 and 0.55 for live weight at slaughter, weight after bleeding and weight with viscera and without head, were ranged from 0.28 to 0.60 for edible parts, were estimates ranged from 0.38 to 0.42 for dressing yield and ranged from 0.42 to 0.55 inedible parts. All possible genetic correlations between carcass characteristics were moderate to high and positive except among TED and both of LU and FUR, also between DR1 and VIS were negative. Common litter correlations among records of carcass characteristics were mostly moderate or high, positive and negative. Correlations of environmental between carcass characteristics were mostly moderate or high, positive and negative. Estimates of ( $r_P$ ) between records of different carcass characteristics were mainly positive and moderate or high magnitude. No significant differences of breed on carcass characteristics except for gible. New Zealand White had highest of gible and leaver. In mostly

parity affected significantly on carcass characteristics, the highest value during first parity except for gible and leaver were highest value during second parity. Contrary other traits were insignificantly. Most carcass characteristics were higher in winter months than other seasons.

**Key words:** Edible parts, dressing yield %, inedible parts, heritability, genetic and phenotypic correlations.

### المستخلص :

تم تحليل البيانات باستخدام برنامج النموذج الحيواني بطريقة معظمة الاحتمال المعتمدة على حساب المشتقات التفاضلية DFREML لخصائص الذبيحة لتقدير المكافىء الوراثي و التأثير المشترك لخلفة البطن و تأثير الخطأ التجريبي و تقدير الارتباط الوراثي والمشارك لخلفة البطن و البيئي والظاهري. تم استخدام البرنامج الإحصائي (SAS 2003) للحصول على المربعات الصغرى لتأثير السلالة والبطن والموسم والمعنوية. بوجه عام كانت قيم المكافىء الوراثي متوسطة وكانت ٠,٢٠ - ٠,١٢ و ٠,٣٧ للوزن الحى عند الذبح وبعد الذبح وبالأحشاء وبدون الرأس. وكانت قيمته متوسطة وعالية وتتراوح بين ٠,٢٩ - ٠,٥٤ للأجزاء الصالحة للأكل. وكانت متوسطة وتتراوح بين ٠,٣٠ - ٠,٣٥ لنسب التصافى. وايضا كانت منخفضة ومتوسطة و تراوحت بين ٠,٠٣ - ٠,٣٣ بالنسبة للأجزاء غير الصالحة للأكل. وكانت قيم التأثير المشترك لخلفة البطن ٠,٣٧ و ٠,٥٥ و ٠,٥٥ للوزن الحى عند الذبح وبعد الذبح وبالأحشاء وبدون الرأس . وتتراوح بين ٠,٣٨ - ٠,٤٢ لنسب التصافى . وتتراوح بين ٠,٤٢ - ٠,٥٥ للأجزاء الصالحة للأكل. جميع الارتباطات الوراثية الممكنة بين خصائص الذبيحة كانت متوسطة وعالية ووموجبة ماعدا بين كلا من الأجزاء الصالحة للأكل و الفرو والرئة. وايضا نسبة التصافى ١ و الاحشاء. والارتباط المشترك لخلفة البطن بين خصائص الذبيحة كانت متوسطة وعالية وموجبة وسالبة. كانت قيم الارتباط البيئي لخصائص الذبيحة كانت متوسطة وعالية وموجبة وسالبة. الارتباط الظاهري بين خصائص الذبيحة المختلفة كان موجب ومتوسط وعالى. لا توجد فروق معنوية لتأثير السلالة على خصائص الذبيحة باستثناء gible و كانت سلالة النيوزيلندى الأبيض أعلى لقيم gible والكبد. كان تأثير البطن بوجه عام معنوى على خصائص الذبيحة وكانت أعلى قيم خلال البطن الأولى ما عدا gible والكبد كانت أعلى خلال البطن الثانية عن باقى الصفات الأخرى. كانت معظم خصائص الذبيحة أعلى فى الشتاء عن المواسم الأخرى.

**التوصية:** من القيم المتوسطة او العالية للمكافء الوراثى فانه يمكن تحسين خصائص الذبيحة من خلال الأنتخاب للحيوانات. كان التأثير المشترك لخلفة البطن واضحا وله تأثيرات قوية على خصائص الذبيحة. كانت قيم الارتباط الوراثي والمشارك لخلفة البطن و البيئى لخصائص الذبيحة متوسط وموجب وعالى فإن الأنتخاب لأحدى الصفات سيؤدى الى التحسين فى الأخرى. لا توجد فروق معنوية لتأثير السلالة على خصائص الذبيحة. كان تأثير البطن بوجه عام معنوى على خصائص الذبيحة. كانت معظم خصائص الذبيحة أعلى فى الشتاء عن المواسم الأخرى.

## INTRODUCTION:

Production of rabbit has an important role in bridging shortage of food in many countries Khalil et al.,(2016). For meat of Rabbit has several advantages including high protein content and low cholesterol content Hanaa et al.,(2014). The quality attributes of food products including rabbit meat have been attracting an increasing interest in recent years. Meat products from ecological rearing were characterized by a higher nutritive quality and better taste Horsted et al.,(2010). In addition rabbit has a quite high dressing percentage when compared to ruminants, ranging between 50 - 65% (Lebas et al., 1986 and Roiron et al., 1992). Heritability of different carcass traits is medium to high, and therefore carcass traits might be considered in rabbit selection and breeding, however it has been reported that different factors, such as age at slaughter, weight at slaughter, breed, and sex, have an influence on different carcass traits (Parigi Bini et al., 1992; Bianospino et al., 2006; Gašperlin et al., 2006 and Metzger et al., 2006). The purpose of this study is to estimate of heritability, common litter effects, and genetic, common litter, environmental and phenotypic using multi-trait animal to determine the suitability of selection for achieving genetic improvement. Also discuss some non-genetic that affect carcass characteristics for New Zealand white and Baladi Black rabbits. Hence, an effort was made to find the effects of different

factors on the carcass characteristics for obtaining maximum dressed meat and dressing yield.

## **MATERIALS AND METHODS:**

### **Animals and data:**

This study was carried out at Sakha Animal Farm, Animal Production Research Institute, Agricultural Research Center, Dokki, Egypt (APRI). Data collected on 120 random male at 12 weeks of age, from New Zealand white and Baladi Black rabbit.

### **Slaughtering procedure:**

All rabbits were weighed and slaughtered. Each rabbit was bled and weighed to determine the blood weight. Fur was removed and weighed after slaughter. Head was separated and the internal organs were also removed and weighed. Fur with tail and feet and other inedible parts were also measured with a sensitive scale. Data collection as followed; Pre-slaughter weight (g) was the live weight of each rabbit before slaughter in grammars; Hot carcass (g) was the weight after slaughter and bleeding of rabbit; Giblets (g) were total weight of kidney, liver and heart; Inedible carcass (g) were total weight of lung, fur with tail and feet, viscera, blood, ; Edible carcass (g, dressed head and gible. Dressing yield (%) was taken as the percentage of edible carcass to the pre-slaughter weight.

### **The statistical analysis:**

The variances and covariances were obtained using REML method of VARCOMP procedure of SAS 2003. Data were analyzed applying multi-trait animal model of carcass characteristics applying MTDFREML programs of Boldman *et al.*, 1995, to obtain

Starting mixed model was obtained applying REML method of VARCOMP procedure of SAS 2003. Data were analyzed using multi-trait animal model of carcass characteristics using MTDFREML programs of Boldman *et al.*, 1995, to obtain

proportion of the phenotypic variance due to additive genetic effects ( $h^2$ ); common litter effects ( $c^2$ ); random error effects ( $e^2$ ); genetic ( $r_G$ ), common litter ( $r_C$ ), environmental ( $r_E$ ) and phenotypic ( $r_P$ ) correlations. Analyses were done according to the general model:

$$y = Xb + Z_1a + Z_2p + e. \quad (\text{Model 1})$$

Where,  $y$ =Vector of observation,  $X$ = Incidence matrix of fixed effects;  $b$  = vector of fixed effects including breed, (NZW and BB), parity (3 levels) and season (4 levels);  $Z_1$  and  $Z_2$ = incidence matrices corresponding to random effects of additive ( $a$ ) and common litter effect ( $r_C$ ) respectively.

Data were carried out by weighed least-squares means method in the procedure GLM of statistical software SAS(2003) to obtain least squares and used to compare means by Duncan's multiple range test for carcass characteristics.

$$Y_{ijk} = \mu + B_i + P_j + S_k + e_{ijk}. \quad (\text{Model 2}).$$

Where:

$Y_{ijk}$  = the parameters on the  $ijk^{\text{th}}$  carcass characteristics,  $\mu$  = the overall mean,  $B_i$  = the fixed effect of the  $i^{\text{th}}$  breeds ( $i$ = NZW and BB);  $P_j$  = the fixed effect of the  $j^{\text{th}}$  parity ( $j$ =1, 2 and 3);  $S_k$  = the fixed effect of the  $k^{\text{th}}$  season ( $k$ =1,2, 3 and 4) and  $e_{ijk}$  = the random deviation of all the other effects not specified in the model.

### III. Results and Discussion:

#### *Genetic affect:*

#### *Heritability estimates*

Heritability estimates for carcass characteristics in (Table 1). The estimates were mostly moderate 0.20, 0.12 and 0.37 for live weight at slaughter, weight after bleeding and weight with viscera and without head. The estimates were moderate and high and ranged from 0.29 to 0.54 for edible parts. The estimates were moderate and ranged from 0.30 to 0.35 for dressing yield. The estimates were low and moderate and ranged from 0.03 to 0.33 for inedible parts. Ayyat *et al.*, (1994) with NZW rabbits,

Heritability were low to moderate for non-edible carcass traits. Ferraz and Eler, (1996) found moderate estimates of heritability for carcass weight and carcass yield of 0.178 and 0.152 for the Californian breed and 0.152 and 0.000 for NZW rabbits, respectively. Khalil et al., (2005) reported heritability estimates for hot carcass weight, offal weight and meat weight were moderate and ranged 0.10 to 0.16 but low for dressing percent (0.097). Al Seaf et al., (2007) reported heritability estimates for edible and non-edible carcass were mostly moderate and ranged from 0.12 to 0.22. Garreau et al., (2008) reported that heritability estimates for carcass yield were moderate (0.24).

Table 1: Estimates for proportion of the phenotypic variance due to additive genetic effects ( $h^2$ ) and to common litter effects ( $c^2$ ) and to random error effects ( $e^2$ ) with standard errors ( $\pm$ SE) for carcass characteristics for New Zealand white (NZW) and Baladi Black (BB) rabbits.

| <b>Arameters</b>                        | <b><math>h^2</math></b> | <b><math>c^2</math></b> | <b><math>e^2</math></b> |
|---|-------------------------|-------------------------|-------------------------|
| Live weight, at slaughter, g            | 0.20 $\pm$ 0.03         | 0.37 $\pm$ 0.06         | 0.43 $\pm$ 0.09         |
| Weight after bleeding, g                | 0.12 $\pm$ 0.02         | 0.55 $\pm$ 0.07         | 0.33 $\pm$ 0.05         |
| Weight with viscera and without head, g | 0.37 $\pm$ 0.06         | 0.54 $\pm$ 0.08         | 0.08 $\pm$ 0.1          |
| <b>Edible parts:</b>                    |                         |                         |                         |
| Hot carcass without head, g             | 0.29 $\pm$ 0.11         | 0.60 $\pm$ 0.09         | 0.11 $\pm$ 0.02         |
| Giblet, g                               | 0.41 $\pm$ 0.06         | 0.52 $\pm$ 0.08         | 0.07 $\pm$ 0.14         |
| Head, g                                 | 0.38 $\pm$ 0.10         | 0.53 $\pm$ 0.08         | 0.11 $\pm$ 0.02         |
| Total edible, g                         | 0.54 $\pm$ 0.07         | 0.28 $\pm$ 0.04         | 0.17 $\pm$ 0.03         |
| <b>Dressing yield %:</b>                |                         |                         |                         |
| Carcass %                               | 0.31 $\pm$ 0.11         | 0.42 $\pm$ 0.06         | 0.27 $\pm$ 0.04         |
| Carcass with giblet %                   | 0.30 $\pm$ 0.11         | 0.39 $\pm$ 0.06         | 0.30 $\pm$ 0.05         |
| Carcass with giblet and head %          | 0.35 $\pm$ 0.10         | 0.38 $\pm$ 0.06         | 0.28 $\pm$ 0.04         |
| <b>Inedible parts:</b>                  |                         |                         |                         |
| Lung, g                                 | 0.16 $\pm$ 0.03         | 0.47 $\pm$ 0.08         | 0.37 $\pm$ 0.06         |
| Fur, g                                  | 0.33 $\pm$ 0.05         | 0.42 $\pm$ 0.02         | 0.25 $\pm$ 0.12         |
| Viscera, g                              | 0.06 $\pm$ 0.15         | 0.55 $\pm$ 0.09         | 0.39 $\pm$ 0.06         |
| Blood, g                                | 0.03 $\pm$ 0.01         | 0.48 $\pm$ 0.08         | 0.49 $\pm$ 0.08         |
| Total inedible                          | 0.11 $\pm$ 0.14         | 0.45 $\pm$ 0.07         | 0.44 $\pm$ 0.07         |
| Inedible : edible                       | 0.19 $\pm$ 0.03         | 0.50 $\pm$ 0.08         | 0.32 $\pm$ 0.11         |

From these observations for moderate or high for hertabilities, it is genetic point of view improvements were achieved in carcass traits through selection of animals.

***Common litter effects***

Common litter effects for carcass characteristics in (Table 1) were moderate to high and were generally higher than the respective heritabilities. Estimates were 0.37, 0.55 and 0.55 for live weight at slaughter, weight after bleeding and weight with

viscera and without head. The estimates ranged from 0.28 to 0.60 for edible parts. The estimates ranged from 0.38 to 0.42 for dressing yield. The estimates ranged from 0.42 to 0.55 for inedible parts. These results are in agreement with Al Seaf et al., (2007) who reported common litter effects for carcass traits and ranged from 0.31 to 0.37 for slaughter and edible carcass traits, 0.29 to 0.39 for non-edible carcass traits and suggested common litter effects appeared to have strong effects on growth even up to slaughtering time. Ferraz et al., (1992) found that common environmental effects to be consistently more important than direct genetic effects for several traits studied, but Lukefahr et al., (1996) reported that for each carcass trait investigated, the magnitudes of variance components for direct genetic and common environmental effects were similar.

#### ***Correlations among carcass characteristics:***

Importance of economic and biological relationships between studying traits may be understood in relationship to other traits. The estimates of genetic ( $r_G$ ), common litter ( $r_C$ ), environmental ( $r_E$ ) and phenotypic ( $r_P$ ) correlations are shown in Tables 2, 3, 4 and 5.

#### ***Genetic correlations ( $r_G$ )***

Generally all possible genetic correlations between carcass characteristics were moderate to high and positive except among TED and both of LU and FUR, also between DR1 and VIS were negative (Table 2). The higher correlations between LWB and both of WB; DR1; LU and IND: TED, also higher among WB and both of WBV; DR1; DR2, also higher among HC and TE, also higher among G and both of TE; VIS, also higher among H and both of DR1; BL, also higher among TED and both of B1; TIND; also higher among DR1; DR2; DR3; BL; TIND, also higher among DR2 and both of DR3; FUR; TIND; IND: TED and higher between TIND and IND: TED. From results

selection improving LWB might be expected to improvement in all pervious traits.Su et al., (1999) found that a negative genetic correlation between post-weaning daily gain and dressing yield (-0.22). Garreau et al., (2008) reported that the negative genetic correlation between 63-day body weight and dressing yield (-0.24). Hanaa et al., (2014) reported that the genetic correlation between slaughter weight and weaning weight were high and positive (0.73).

Table 2: Estimates genetic correlations ( $r_G$ ) for carcass characteristics for (NZW) and (BB) rabbits.

|             | WB   | WBV  | HC   | G    | H    | TED  | DR1  | DR2  | DR3  | LU    | FUR   | VIS   | BL   | TIND | TIND:TED |
|-------------|------|------|------|------|------|------|------|------|------|-------|-------|-------|------|------|----------|
| <b>LWB</b>  | 0.96 | 0.56 | 0.59 | 0.57 | 0.49 | 0.19 | 0.98 | 0.54 | 0.53 | 0.86  | 0.57  | 0.49  | 0.04 | 0.78 | 0.97     |
| <b>WB</b>   |      | 0.86 | 0.76 | 0.58 | 0.44 | 0.53 | 0.88 | 0.81 | 0.62 | 0.59  | 0.51  | 0.68  | 0.62 | 0.14 | 0.78     |
| <b>WBV</b>  |      |      | 0.60 | 0.51 | 0.79 | 0.81 | 0.54 | 0.40 | 0.50 | 0.32  | 0.36  | 0.64  | 0.31 | 0.93 | 0.42     |
| <b>HC</b>   |      |      |      | 0.29 | 0.34 | 0.99 | 0.62 | 0.50 | 0.50 | 0.58  | 0.68  | 0.64  | 0.42 | 0.54 | 0.46     |
| <b>G</b>    |      |      |      |      | 0.91 | 0.98 | 0.45 | 0.61 | 0.54 | 0.76  | 0.36  | 0.91  | 0.72 | 0.39 | 0.57     |
| <b>H</b>    |      |      |      |      |      | 0.47 | 0.85 | 0.28 | 0.21 | 0.47  | 0.33  | 0.60  | 0.85 | 0.65 | 0.64     |
| <b>TED</b>  |      |      |      |      |      |      | 0.20 | 0.33 | 0.26 | -0.73 | -0.63 | 0.66  | 0.82 | 0.88 | 0.74     |
| <b>DR1</b>  |      |      |      |      |      |      |      | 0.99 | 0.99 | 0.57  | 0.17  | -0.29 | 0.89 | 0.66 | 0.80     |
| <b>DR2</b>  |      |      |      |      |      |      |      |      | 0.99 | 0.77  | 0.98  | 0.82  | 0.93 | 0.99 | 0.99     |
| <b>DR3</b>  |      |      |      |      |      |      |      |      |      | 0.91  | 0.88  | 0.94  | 0.87 | 0.46 | 0.81     |
| <b>LU</b>   |      |      |      |      |      |      |      |      |      |       | 0.67  | 0.57  | 0.43 | 0.79 | 0.77     |
| <b>FUR</b>  |      |      |      |      |      |      |      |      |      |       |       | 0.82  | 0.78 | 0.71 | 0.15     |
| <b>VIS</b>  |      |      |      |      |      |      |      |      |      |       |       |       | 0.73 | 0.85 | 0.87     |
| <b>BL</b>   |      |      |      |      |      |      |      |      |      |       |       |       |      | 0.61 | 0.70     |
| <b>TIND</b> |      |      |      |      |      |      |      |      |      |       |       |       |      |      | 0.97     |

LWB=Live weight, at slaughter, g; WB=Weight after bleeding, g; WBV=Weight with viscera and without head, g; HC=Hot carcass without head, g; G=Giblet, g; H=Head, g; TED= Total edible, g; DR1= Carcass %; DR2= Carcass with giblet %; DR3= Carcass with giblet and head %; LU= Lung, g; VIS= Viscera, g; B=Blood; TIND= Total inedible; IND:TED= Inedible : edible.

***Common litter correlations ( $r_C$ )***

The estimates of common litter correlations among records of carcass characteristics mostly moderate or high, positive and negative (Table 3). The higher correlations among LWB and both of and both of WBV; HC; H; TED ; DR2; DR3 and IND: TED, also higher among WB and both of WBV; FUR and TIND, also

higher among HC and TE, also higher among G and both of H TED, also higher among DR1and both of DR2; DR3; LU; BL.also higher among DR2and both of B1; TIND; also higher among DR1; DR2; DR3; BL; TIND, also higher among DR2 and both of DR3; FUR; VIS TIND, higher amongDR3 and both ofLU; FUR; VIS, higher between LU and VIS and higher between TIND and IND: TED. From resultsthe importance ofcommon litter correlationoncarcasscharacteristics becomes evident.Thus, it is necessary might recommend that these correlations should be considered in any program of breeding to improve litter traits in rabbits.No research is available for estimates of( $r_C$ ) carcass characteristics in rabbits.

Table 3: Estimates common litter correlations ( $r_C$ ) for carcass characteristics for (NZW) and (BB) rabbits.

|      | WB   | WBV  | HC   | G    | H    | TED  | DR1   | DR2   | DR3   | LU    | FUR   | VIS   | BL    | TIND  | TIND:TED |
|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|----------|
| LWB  | 0.72 | 0.92 | 0.93 | 0.66 | 0.94 | 0.98 | 0.41  | 0.98  | 0.80  | 0.60  | 0.64  | 0.49  | 0.61  | 0.76  | 0.96     |
| WB   |      | 0.97 | 0.50 | 0.53 | 0.70 | 0.79 | -0.89 | -0.56 | 0.63  | 0.50  | 0.92  | 0.68  | -0.11 | 0.82  | 0.17     |
| WBV  |      |      | 0.67 | 0.28 | 0.27 | 0.73 | -0.99 | -0.98 | -0.97 | 0.54  | 0.43  | 0.62  | 0.55  | 0.58  | 0.19     |
| HC   |      |      |      | 0.17 | 0.22 | 0.98 | 0.58  | 0.60  | 0.51  | 0.31  | -0.57 | -0.96 | 0.13  | -0.78 | -0.90    |
| G    |      |      |      |      | 0.93 | 0.97 | -0.89 | -0.96 | -0.71 | -0.98 | 0.34  | -0.13 | -0.72 | 0.20  | -0.37    |
| H    |      |      |      |      |      | 0.98 | -0.97 | -0.98 | -0.84 | -0.18 | 0.38  | 0.21  | -0.13 | 0.25  | -0.45    |
| TED  |      |      |      |      |      |      | 0.76  | 0.54  | 0.56  | 0.63  | 0.76  | 0.41  | -0.97 | 0.74  | 0.67     |
| DR1  |      |      |      |      |      |      |       | 0.98  | 0.99  | 0.94  | -0.98 | -0.89 | 0.98  | 0.77  | 0.48     |
| DR2  |      |      |      |      |      |      |       |       | 0.98  | 0.50  | 0.80  | 0.90  | -0.34 | 0.99  | -0.99    |
| DR3  |      |      |      |      |      |      |       |       |       | 0.91  | 0.91  | 0.88  | -0.50 | 0.42  | 0.35     |
| LU   |      |      |      |      |      |      |       |       |       |       | 0.75  | 0.81  | 0.77  | 0.11  | 0.78     |
| FUR  |      |      |      |      |      |      |       |       |       |       |       | 0.29  | -0.17 | 0.43  | 0.77     |
| VIS  |      |      |      |      |      |      |       |       |       |       |       |       | 0.49  | 0.92  | 0.88     |
| BL   |      |      |      |      |      |      |       |       |       |       |       |       |       | 0.67  | 0.13     |
| TIND |      |      |      |      |      |      |       |       |       |       |       |       |       |       | 0.96     |

+ Trait as defined in Table 2.

### *Environmental correlations ( $r_E$ )*

All possible correlations of environmental correlations between carcass characteristics mostly moderate or high, positiveand negative (Table 4).Falconer and Mackay (1989)they reported some cases, estimates of  $r_G$  and  $r_E$ are different in magnitude, or even in sign, while in other cases the two types of correlations are of the same sign and do not differ in magnitude,and a large difference, and particularly a difference in sign, showed that genetic and environmental sources of variation

affect the characters through different physiological mechanism and this is most common mode of my study. No research is available for estimates of ( $r_E$ ) carcass characteristics in rabbits.

Table 4: Estimates environmental correlations ( $r_E$ ) for carcass characteristics for (NZW) and (BB) rabbits.

|      | WB   | WBV  | HC   | G    | H    | TED  | DR1   | DR2   | DR3   | LU    | FUR   | VIS   | BL    | TIND | TIND:TED |
|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|------|----------|
| LWB  | 0.96 | 0.91 | 0.38 | 0.70 | 0.46 | 0.82 | -0.68 | 0.17  | 0.88  | 0.04  | 0.72  | 0.73  | -0.19 | 0.79 | -0.49    |
| WB   |      | 0.77 | 0.59 | 0.42 | 0.71 | 0.68 | -0.01 | -0.97 | -0.94 | 0.28  | 0.37  | 0.53  | 0.61  | 0.85 | 0.66     |
| WBV  |      |      | 0.64 | 0.44 | 0.80 | 0.55 | 0.78  | 0.53  | 0.31  | 0.33  | 0.40  | 0.91  | 0.76  | 0.98 | 0.44     |
| HC   |      |      |      | 0.63 | 0.58 | 0.88 | 0.53  | 0.50  | 0.50  | 0.18  | 0.91  | 0.45  | 0.32  | 0.54 | 0.99     |
| G    |      |      |      |      | 0.69 | 0.94 | 0.54  | 0.95  | 0.97  | 0.99  | 0.85  | 0.58  | 0.75  | 0.61 | 0.34     |
| H    |      |      |      |      |      | 0.61 | 0.23  | 0.83  | 0.92  | 0.83  | 0.25  | 0.35  | 0.55  | 0.55 | 0.55     |
| TED  |      |      |      |      |      |      | 0.64  | 0.64  | 0.65  | 0.71  | 0.41  | -0.78 | 0.90  | 0.63 | 0.82     |
| DR1  |      |      |      |      |      |      |       | 0.99  | 0.12  | -0.06 | -0.24 | 0.98  | 0.19  | 0.82 | 0.82     |
| DR2  |      |      |      |      |      |      |       |       | 0.98  | -0.61 | 0.80  | 0.99  | -0.97 | 0.98 | 0.98     |
| DR3  |      |      |      |      |      |      |       |       |       | 0.96  | 0.91  | 0.88  | -0.50 | 0.42 | 0.35     |
| LU   |      |      |      |      |      |      |       |       |       |       | 0.75  | 0.81  | 0.77  | 0.11 | 0.44     |
| FUR  |      |      |      |      |      |      |       |       |       |       |       | 0.74  | 0.48  | 0.83 | 0.64     |
| VIS  |      |      |      |      |      |      |       |       |       |       |       |       | 0.67  | 0.90 | 0.91     |
| BL   |      |      |      |      |      |      |       |       |       |       |       |       |       | 0.18 | 0.61     |
| TIND |      |      |      |      |      |      |       |       |       |       |       |       |       |      | 0.61     |

+ Trait as defined in Table 2.

***Phenotypic correlations ( $r_P$ )***

Estimates of ( $r_P$ ) between records of different carcass characteristics were mainly positive and moderate or high magnitude (Table 5). Hanaa et al., (2014) reported that daily weight gain from weaning to slaughter was phenotypically high positive correlated with slaughter weight with value of 0.68.

However, positive ( $r_P$ ) among any two traits of carcass characteristics do not necessarily indicate that selecting on one of these trait will lead to an improvement in the other, because a phenotypic correlation is not always a reliable estimate of the genetic relationship existing between traits, for example, environmental effect on two traits could be strong and positively correlated that a negative genetic correlation is masked. Therefore, it is recognized that phenotype association is not a satisfactory guide to the expected correlated genetic response of traits under selection; only genetic correlation should be used for such a prediction. In practice, high or moderate and positive estimates of the phenotypic correlation in current studies

and review among carcass characteristics give considerable advantage for rabbit breeders in their management and culling decisions.

Table 5: Estimates phenotypic correlations ( $r_p$ ) for carcass characteristics for (NZW) and (BB) rabbits.

|      | WB   | WBV  | HC   | G    | H    | TED  | DR1   | DR2   | DR3   | LU    | FUR   | VIS   | BL    | TIND | TIND:TED |
|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|------|----------|
| LWB  | 0.83 | 0.74 | 0.66 | 0.57 | 0.65 | 0.11 | 0.37  | 0.73  | 0.40  | 0.62  | 0.58  | 0.58  | 0.18  | 0.77 | 0.72     |
| WB   |      | 0.58 | 0.52 | 0.48 | 0.42 | 0.61 | -0.26 | -0.64 | 0.65  | 0.41  | 0.60  | 0.61  | 0.28  | 0.77 | 0.42     |
| WBV  |      |      | 0.64 | 0.49 | 0.50 | 0.72 | -0.24 | -0.33 | -0.32 | 0.43  | 0.39  | 0.63  | 0.42  | 0.67 | 0.26     |
| HC   |      |      |      | 0.25 | 0.31 | 0.97 | 0.57  | 0.51  | 0.48  | 0.33  | 0.08  | -0.38 | 0.15  | 0.18 | 0.11     |
| G    |      |      |      |      | 0.91 | 0.98 | -0.17 | -0.08 | 0.03  | 0.02  | 0.40  | 0.17  | -0.15 | 0.29 | 0.19     |
| H    |      |      |      |      |      | 0.55 | -0.12 | -0.20 | -0.14 | 0.20  | 0.33  | 0.27  | 0.15  | 0.37 | 0.15     |
| TED  |      |      |      |      |      |      | 0.48  | 0.46  | 0.43  | 0.19  | 0.08  | 0.09  | 0.08  | 0.62 | 0.67     |
| DR1  |      |      |      |      |      |      |       | 0.98  | 0.98  | 0.38  | -0.46 | -0.80 | 0.84  | 0.52 | 0.65     |
| DR2  |      |      |      |      |      |      |       |       | 0.95  | 0.99  | -0.15 | 0.70  | -0.15 | 0.95 | 0.97     |
| DR3  |      |      |      |      |      |      |       |       |       | -0.13 | 0.85  | 0.98  | -0.13 | 0.49 | 0.31     |
| LU   |      |      |      |      |      |      |       |       |       |       | 0.27  | 0.61  | 0.23  | 0.44 | 0.51     |
| FUR  |      |      |      |      |      |      |       |       |       |       |       | 0.47  | 0.17  | 0.74 | 0.56     |
| VIS  |      |      |      |      |      |      |       |       |       |       |       |       | 0.56  | 0.89 | 0.85     |
| BL   |      |      |      |      |      |      |       |       |       |       |       |       |       | 0.42 | 0.34     |
| TIND |      |      |      |      |      |      |       |       |       |       |       |       |       |      | 0.86     |

+ Trait as defined in Table 2

### Non-genetic affect:

#### *Breed effect*

Least square means of the carcass characteristics of New Zealand White (NZW) and Baladi Black (BB) were presented on Table 6 shows that in generally there were no significant differences of breed on carcass characteristics except for gible. New Zealand White had highest of gible and leaver. Ghosh et al. (2004) reported that no significant differences among breeds in body weight at maturity. Ouyed and Brun (2008) found that there were no significant effects of breed type on commercial carcass weight, commercial carcass yield of New Zealand White and Californian breed and their crosses. Ghosh and Mandel (2008) reported that the effects of breed on the values of hot carcass, gible, dressed head and total edible were non-significant. Baiomy and Hassanien (2011) observed that though breed differences had no significant effect on most carcass traits, dressing yield of carcass was significantly higher in New Zealand white than Californian breed (58.5, 57.3%), respectively. While Das and Bardoloi (2008) they reported that breed had

highly significantly ( $P < 0.01$ ) effects on carcass weight of rabbits and inedible offal and breed did not have a significant effect on edible organs. However Fadare (2015) reported that the New Zealand white had the highest dressing yield percentage of  $67.95 \pm 1.95$  while Palomino brown had the least dressing yield ( $55.23 \pm 2.36\%$ ). California and Havana black breed had similar dressing yield.

Table 6. Least-squares means and standard errors of carcass characteristics as affected by breed for New Zealand white (NZW) and Baladi Black (BB) rabbits.

| Parameters                              | NZW               | BB                |
|---|-------------------|-------------------|
| Live weight, at slaughter, g            | $1970.4 \pm 33.5$ | $1956.5 \pm 33.5$ |
| Weight after bleeding, g                | $1926.1 \pm 33.7$ | $1910.6 \pm 33.7$ |
| Weight with viscera and without head, g | $1461.6 \pm 18.9$ | $1024.8 \pm 18.9$ |
| <b>Edible parts:</b>                    |                   |                   |
| Hot carcass without head, g             | $1013.5 \pm 18.9$ | $1024.8 \pm 18.9$ |
| Giblet, g                               | $107.1 \pm 2.9^a$ | $88.6 \pm 2.9^b$  |
| Head, g                                 | $115.8 \pm 2.1$   | $119.4 \pm 2.1$   |
| Total edible, g                         | $1236.4 \pm 20.5$ | $1232.9 \pm 20.5$ |
| <b>Dressing yield %:</b>                |                   |                   |
| Carcass %                               | $51.9 \pm 0.90$   | $52.5 \pm 0.90$   |
| Carcass with giblet %                   | $57.3 \pm 0.91$   | $57.1 \pm 0.91$   |
| with giblet and head %                  | $58.9 \pm 0.92$   | $59.7 \pm 0.92$   |
| <b>Inedible parts:</b>                  |                   |                   |
| Lung, g                                 | $12.2 \pm 0.41$   | $13.3 \pm 0.41$   |
| Fur, g                                  | $348.5 \pm 11.6$  | $321.1 \pm 11.6$  |
| Viscera, g                              | $452.0 \pm 19.0$  | $448.0 \pm 19.0$  |
| Blood, g                                | $40.6 \pm 2.3$    | $41.9 \pm 2.3$    |
| Total inedible                          | $853.2 \pm 27.6$  | $824.2 \pm 27.6$  |
| Inedible : edible                       | $69.1 \pm 2.3$    | $68.0 \pm 2.3$    |

### **Parity effect:**

In mostly parity affected significantly ( $P \leq 0.01$ ) (LBW, SBW, SBWV, SBWN, hot carcass, giblet, total edible, kidney and leaver), the highest value during first parity except for giblet

and leaver were highest value during second parity. Contrary other traits were insignificantly (Table 7). Whereas Prayaga and Eady (2003). They obtained significantly higher carcass weight in 2<sup>nd</sup> and 3<sup>rd</sup> parity litters than in 1<sup>st</sup> and 4<sup>th</sup> ones. Ouyed and Brun (2008) reported that there was no significant effect of parity on carcass traits.

Table 7. Least-squares means and standard errors of carcass characteristics as affected by parity for (NZW) and (BB) rabbits.

| Parameters                              | 1 <sup>st</sup>            | 2 <sup>nd</sup>                         | 3 <sup>rd</sup>            |
|---|----------------------------|---|----------------------------|
| Live weight, at slaughter, g            | 2011.3 ± 60.8 <sup>a</sup> | 1981.2 ± 60.8 <sup>ab</sup>             | 1936.3 ± 47.1 <sup>b</sup> |
| Weight after bleeding, g                | 1975.8 ± 61.2 <sup>a</sup> | 1926.9 ± 61.2 <sup>ab</sup>             | 1893.1 ± 47.4 <sup>b</sup> |
| Weight with viscera and without head, g | 1541.6 ± 46.3 <sup>a</sup> | 1444.0 ± 46.3 <sup>a</sup> <sup>b</sup> | 1437.6 ± 35.9 <sup>b</sup> |
| <b>Edible parts:</b>                    |                            |   |                            |
| Hot carcass without head, g             | 1071.3 ± 36.1 <sup>a</sup> | 989.1 ± 36.7 <sup>b</sup>               | 988.3 ± 27.9 <sup>b</sup>  |
| Giblet, g                               | 91.3 ± 4.7 <sup>a</sup>    | 109.4 ± 4.7 <sup>ab</sup>               | 105.3 ± 3.6 <sup>b</sup>   |
| Head, g                                 | 114.1 ± 3.8                | 122.6 ± 3.8                             | 115.6 ± 3.0                |
| Total edible, g                         | 1276.6 ± 39.0 <sup>a</sup> | 1221.1 ± 39.0 <sup>b</sup>              | 1209.2 ± 30.2 <sup>b</sup> |
| <b>Dressing yield %:</b>                |                            |   |                            |
| Carcass %                               | 53.4 ± 1.6                 | 50.5 ± 1.6                              | 51.3 ± 1.3                 |
| Carcass with giblet %                   | 57.9 ± 1.7                 | 56.0 ± 1.7                              | 56.8 ± 1.3                 |
| Carcass with giblet and head %          | 60.0 ± 1.7                 | 57.8 ± 1.7                              | 58.4 ± 1.3                 |
| <b>Inedible parts:</b>                  |                            |   |                            |
| Lung, g                                 | 14.2 ± 0.98                | 11.4 ± 0.78                             | 12.3 ± 0.60                |
| Fur, g                                  | 317.2 ± 34.3               | 375.8 ± 21.3                            | 330.4 ± 16.5               |
| Viscera, g                              | 473.0 ± 34.3               | 467.7 ± 34.3                            | 432.0 ± 26.6               |
| Blood, g                                | 45.4 ± 4.6                 | 38.1 ± 4.6                              | 41.1 ± 3.5                 |
| Total inedible                          | 843.1 ± 48.5               | 886.3 ± 48.5                            | 835.6 ± 37.5               |
| Inedible : edible                       | 68.4 ± 4.2                 | 71.3 ± 4.2                              | 69.2 ± 3.3                 |

### *Season effect:*

In generally the effect of season on carcass characteristics in (Table 8) was higher significantly ( $P \leq 0.01$ ). Most carcass characteristics were higher in winter months than other seasons, may be attributed to suitability of the environmental conditions to rabbit production, increase in food consumption and abundance of

green fodders during winter months and summer is more stressful due to extreme heat and relative humidity. Farghaly and El-Mahdy (1999), they reported that season appears to be the major non genetic factor affect live body weight and carcass traits. The highest live body weight and carcass traits were estimated during winter. Antonella (2000) reported that the involved mostly the environmental and the season in which temperature play a major role on productive and slaughtering performance. As occurs for livestock, even for rabbits, the increase of environmental temperature over the thermo neutrality value reduces the feed intake and consequently, growth rate resulting in lower slaughter weight, at commerciality slaughter age, and sometimes, better slaughter yield because of the lower proportion of skin, empty cut and offal's.

Table 8. Least-squares means and standard errors of carcass characteristics as affected by season for (NZW) and (BB) rabbits.

| Parameters                              | 1 <sup>st</sup>            | 2 <sup>nd</sup>            | 3 <sup>rd</sup>             | 4 <sup>th</sup>             |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|
| Live weight, at slaughter, g            | 2049.0 ± 79.0 <sup>a</sup> | 1930.4 ± 79.8 <sup>b</sup> | 1882.0 ± 48.0 <sup>b</sup>  | 1995.8 ± 48.0 <sup>b</sup>  |
| Weight after bleeding, g                | 2046.5 ± 79.0 <sup>a</sup> | 1898.7 ± 79.6 <sup>b</sup> | 1833.2 ± 48.3 <sup>b</sup>  | 1949.4 ± 48.3 <sup>b</sup>  |
| Weight with viscera and without head, g | 1505.9 ± 46.9              | 1019.4 ± 46.9              | 1024.4 ± 28.4               | 1024.4 ± 28.4               |
| <b>Edible parts:</b>                    |                            |                            |                             |                             |
| Hot carcass without head, g             | 995.4 ± 46.9               | 1019.4 ± 46.9              | 1025.6 ± 28.4               | 1024.4 ± 28.4               |
| Giblet, g                               | 137.9 ± 6.1 <sup>a</sup>   | 91.1 ± 6.1 <sup>b</sup>    | 92.2 ± 3.7 <sup>b</sup>     | 86.8 ± 5.7 <sup>b</sup>     |
| Head, g                                 | 126.0 ± 5.0 <sup>a</sup>   | 107.5 ± 5.0 <sup>b</sup>   | 114.0 ± 3.0 <sup>b</sup>    | 122.2 ± 3.0 <sup>a</sup>    |
| Total edible, g                         | 1259.2 ± 50.6 <sup>a</sup> | 1218.0 ± 50.6 <sup>b</sup> | 1231.9 ± 30.7 <sup>ab</sup> | 1233.4 ± 30.7 <sup>ab</sup> |
| <b>Dressing yield %:</b>                |                            |                            |                             |                             |
| Carcass %                               | 48.1 ± 2.1 <sup>b</sup>    | 52.7 ± 2.1 <sup>b</sup>    | 54.8 ± 1.3 <sup>a</sup>     | 51.4 ± 1.3 <sup>b</sup>     |
| Carcass with giblet %                   | 54.7 ± 2.2 <sup>b</sup>    | 57.5 ± 2.2 <sup>b</sup>    | 59.8 ± 1.3 <sup>a</sup>     | 55.7 ± 1.3 <sup>b</sup>     |
| Carcass with giblet and head %          | 55.3 ± 2.2 <sup>b</sup>    | 59.4 ± 2.2 <sup>b</sup>    | 61.9 ± 1.3 <sup>a</sup>     | 58.6 ± 1.3 <sup>b</sup>     |
| <b>Inedible parts:</b>                  |                            |                            |                             |                             |
| Lung, g                                 | 9.9 ± 1.0 <sup>b</sup>     | 14.6 ± 1.0 <sup>a</sup>    | 12.1 ± 0.61 <sup>ab</sup>   | 13.8 ± 0.61 <sup>ab</sup>   |
| Fur, g                                  | 418.6 ± 27.6 <sup>a</sup>  | 301.7 ± 27.6 <sup>b</sup>  | 325.2 ± 16.8 <sup>b</sup>   | 319.0 ± 16.8 <sup>b</sup>   |
| Viscera, g                              | 518.2 ± 44.6 <sup>a</sup>  | 442.2 ± 44.6 <sup>b</sup>  | 395.6 ± 27.1 <sup>b</sup>   | 474.2 ± 27.1 <sup>ab</sup>  |
| Blood, g                                | 39.8 ± 6.0                 | 45.1 ± 6.0                 | 36.8 ± 3.6                  | 44.4 ± 3.6                  |
| Total inedible                          | 995.3 ± 63.0 <sup>a</sup>  | 812.4 ± 63.0 <sup>b</sup>  | 751.9 ± 38.2 <sup>c</sup>   | 860.3 ± 38.2 <sup>b</sup>   |
| Inedible : edible                       | 76.7 ± 5.5 <sup>a</sup>    | 69.2 ± 5.5 <sup>a</sup>    | 61.3 ± 3.3 <sup>b</sup>     | 71.3 ± 3.3 <sup>a</sup>     |

**CONCLUSION:**

From the values of the moderate or high for heritabilities, it is genetic point of view improvements were achieved in carcass traits through selection of animals. Common litter effects

appeared to have strong effects on carcass characteristics. Genetic, common litter, environmental correlations among records of carcass characteristics were mostly moderate or high, positive indicate that selecting on one of these trait will lead to an improvement in the other. No significant differences of breed on carcass characteristics. Mostly parity affected significantly on carcass characteristics. Most carcass characteristics were higher in winter months than other seasons

**REFERENCES:**

- Adelodun Opeyemi Fadare (2015).** Carcass Traits of New Zealand White, Californian, Palomino Brown and Havana Black Rabbit In the Humid Tropics *Journal of Agriculture and Veterinary Science* Volume 8, Issue 3 Ver. I, PP 19-23.
- Al-Saef, A.M., Khalil, M.H., Al-Homidan, A.H., Al-Dobaib, S.N., Al-Sobayil, K.A.(2007).** Carcass, lean composition and meat quality traits in new Saudi synthetic lines of rabbits developed to be adaptable to hot climates. 3rd International Conference of Quantitative Genetics, Zhejiang University, Hangzhou, China, August 19-2.
- Antonella Dallae Zotte (2000).** Main factors influencing the rabbit carcass and meat quality *Conference: 7th World Rabbit Congress, Volume: A*
- Ayyat, M.S., Anous, M.R. and Sadek, M.H. (1994).** Genetic parameters for meat production in rabbits. 1. Non carcass components. *World Rabbit Science*, 2(3): 93-99.
- Baiomy, A.A. and Hassanien, H. H. M.(2011).** Effect of breed and sex on carcass characteristics and meat chemical composition of New Zealand white and Californian rabbits under upper Egyptian environment. *Egyptian Poultry Science* 31 (2): 275-284.
- Bianospino E., Wechsler F.S., Fernandes S., Roça R.O., Moura A.S.A.M.T. (2006).** Growth, carcass and meat quality traits of straight bred and crossbred Botucatu rabbits. *World Rabbit Sci.* 14: 237-246.
- Boldmann, K. G.; Kriese, L. A.; Van Tassell, C. P.; Kachman, S. D. (1995).** A manual for use of MTDFREML. A set programs to obtain estimates of variances and covariance [DRAFT]. U.S. Department of Agriculture, Agricultural Research Service, USA.

- Das S.K. and Bardoloi R.K. (2008).** Study on the factors affecting carcass traits of broiler rabbits in eastern Himalayan region of India *World Rabbit Sci.* 16: 107 – 110.
- Farghaly H M and El-Mahdy M R M (1999).** Genetic and non-genetic factors affecting live, carcass and non-carcass traits of New Zealand White rabbits in Egypt. *Indian Journal of Animal Science* 69(8): 596-603.
- Ferraz, J. B. S.; Johnson, R. K.; Van Vleck, L. D. (1992).** Estimation of genetic trends and genetic parameters for reproductive and growth traits of rabbits raised in subtropics with animal model. *J. Applied Rabbit Research*, 15:131-142.
- Ferraz, J.B.S. and Eler, J.P. (1996).** Comparison of animal models for estimation of (co)variance components and genetic parameters of reproductive, growth and slaughter traits of Californian and New Zealand rabbits raised under tropical conditions. *Proceedings of 6<sup>th</sup> World Rabbit Congress, Vol 2: 279-284. Toulouse, France.*
- Garreau H., Licois D., Rupp R., Rochambeau H. de (2006).** Genetic variability of the resistance to epizootic rabbit enteropathy (ERE): new results. *In Proc. 8<sup>th</sup> World. Cong. Appl. Livest. Prod., Belo Horizonte, Brazil.*
- Garreau H.; Eady S.J.; Hurtaud J., Legarra A. (2008).** Genetic parameters of production traits and resistance to digestive disorders in a commercial rabbit population. *9<sup>th</sup> World Rabbit Congress – June 10-13– Verona– Italy*
- Ghosh N. and Mandal L. (2008).** Carcass and meat quality traits of rabbits (*Oryctolagus cuniculus*) under warm-humid

- condition of West Bengal, India. *Livestock Research for Rural Development* 20 (9)[7].
- Ghosh, S. K.; Das, A.; Bujarbaruah, K. M.; Das, Asit; Dhiman, K. R. and Singh, N. P. (2004).** Studies on the productive and reproductive performances of Soviet Chinchilla and New Zealand White breeds of rabbit under the subtropical condition of Tripura. In: *Proceeding 8th World Rabbit Congress*, 7-10 September, Puebla, Mexico, 1188 -1193.[8].
- Hanaa A, El-Raffa A, Shebl MK, El-Delebshany A, El-Sayed N (2014).** Genetic evaluation of some economic traits in a maternal line of rabbits. *Egyptian Journal of Poultry Science* 34:85-98.
- Horsted K, Allesen-Holm BH, Hermansen JE (2010).** The effect of breed and feed-type on the sensory profile on breast meat in male broilers reared in an organic free-range system. *British Poultry Science* 4:515-524.
- Khalil MH, Shebl MK, Kosba MA, El-Sabrouf K, Zaki N (2016).** Estimate the contribution of incubation parameters influence egg hatchability using multiple linear regression analysis. *Veterinary World* 9:806-810.
- Metzger S., Odermatt M., Szendro Z., Mohaupt M., Romvari R., Makai A, Biro-Nemeth E. , Sipos L. Radnai I., Horn P. (2006).** A study of the carcass traits of different rabbit genotypes. *World Rabbit Science*, 14: 107 -114[9].
- Lebas F, Coudert P, Rouvier R, De Rockambeau H (1986).** *The Rabbit: Husbandry, Health and Production*. F.A.O., U.N, Rome, Italy.
- Lukefahr, S.D., Odi, H.B.and Atakora, J.K.A.(1996).** Mass selection for 70 day body weight in rabbits. *Journal of Animal Science*,74: 1481-1489.

- Ouyed A. and Brun J.M. (2008).** Heterosis, direct and maternal additive effects on rabbit growth and carcass characteristics. In Proc: 9th World Rabbit Congress – June 10-13, 2008 – Verona – Italy 195-200[1].
- Parig-Bini R., Xiccato G., Cinetto M., Dalle Zotte A. (1992).** Effect of age, slaughter weight and sex on carcass and meat quality of rabbit. Anim. Breed Abst., 61: 257.
- Prayaga, K. C. and Eady, S. J. (2003).** Performance of purebred and crossbred rabbits in Australia: Individual growth and slaughter traits. Australian Journal Agriculture Research, 54: 159 – 166.
- Roiron A., Ouhayoun J., Delmas D. (1992).** Effect of body weight and age at slaughter on carcass and meat quality of rabbit. Anim. Breed Abst., 61: 64.
- SAS., (2003).** SAS OnlineDoc 9.13 SAS Institute Inc., Cary, NC, USA.
- Su G., Kjaer J.B., Brenøe U.T., Sørensen P. (1999).** Estimates of genetic parameters in Danish white rabbits using an animal model: I. Growth and carcass traits. World Rabbit Sci., 7, 59–64.

